


**AP<sup>®</sup> CALCULUS AB  
2002 SCORING GUIDELINES (Form B)**

**Question 1**

Let  $R$  be the region bounded by the  $y$ -axis and the graphs of

$y = \frac{x^3}{1+x^2}$  and  $y = 4 - 2x$ , as shown in the figure above.

- (a) Find the area of  $R$ .
- (b) Find the volume of the solid generated when  $R$  is revolved about the  $x$ -axis.
- (c) The region  $R$  is the base of a solid. For this solid, each cross section perpendicular to the  $x$ -axis is a square. Find the volume of this solid.



**Question 2**

The number of gallons,  $P(t)$ , of a pollutant in a lake changes at the rate  $P'(t) = 1 - 3e^{-0.2\sqrt{t}}$  gallons per day, where  $t$  is measured in days. There are 50 gallons of the pollutant in the lake at time  $t = 0$ . The lake is considered to be safe when it contains 40 gallons or less of pollutant.

- (a) Is the amount of pollutant increasing at time  $t = 9$ ? Why or why not?
- (b) For what value of  $t$  will the number of gallons of pollutant be at its minimum? Justify your answer.
- (c) Is the lake safe when the number of gallons of pollutant is at its minimum? Justify your answer.
- (d) An investigator uses the tangent line approximation to  $P(t)$  at  $t = 0$  as a model for the amount of pollutant in the lake. At what time  $t$  does this model predict that the lake becomes safe?

**Question 3**

A particle moves along the  $x$ -axis so that its velocity  $v$  at any time  $t$ , for  $0 \leq t \leq 16$ , is given by  $v(t) = e^{2\sin t} - 1$ . At time  $t = 0$ , the particle is at the origin.


- (a) On the axes provided, sketch the graph of  $v(t)$  for  $0 \leq t \leq 16$ .
- (b) During what intervals of time is the particle moving to the left? Give a reason for your answer.
- (c) Find the total distance traveled by the particle from  $t = 0$  to  $t = 4$ .
- (d) Is there any time  $t$ ,  $0 < t \leq 16$ , at which the particle returns to the origin? Justify your answer.

### Question 4

The graph of a differentiable function  $f$  on the closed interval  $[-3, 15]$  is shown in the figure above. The graph of  $f$  has a horizontal tangent line at  $x = 6$ . Let

$$g(x) = 5 + \int_6^x f(t) dt \text{ for } -3 \leq x \leq 15.$$

- Find  $g(6)$ ,  $g'(6)$ , and  $g''(6)$ .
- On what intervals is  $g$  decreasing? Justify your answer.
- On what intervals is the graph of  $g$  concave down? Justify your answer.
- Find a trapezoidal approximation of  $\int_{-3}^{15} f(t) dt$  using six subintervals of length  $\Delta t = 3$ .



### Question 5

Consider the differential equation  $\frac{dy}{dx} = \frac{3-x}{y}$ .

- Let  $y = f(x)$  be the particular solution to the given differential equation for  $1 < x < 5$  such that the line  $y = -2$  is tangent to the graph of  $f$ . Find the  $x$ -coordinate of the point of tangency, and determine whether  $f$  has a local maximum, local minimum, or neither at this point. Justify your answer.
- Let  $y = g(x)$  be the particular solution to the given differential equation for  $-2 < x < 8$ , with the initial condition  $g(6) = -4$ . Find  $y = g(x)$ .

### Question 6

Ship  $A$  is traveling due west toward Lighthouse Rock at a speed of 15 kilometers per hour (km/hr). Ship  $B$  is traveling due north away from Lighthouse Rock at a speed of 10 km/hr. Let  $x$  be the distance between Ship  $A$  and Lighthouse Rock at time  $t$ , and let  $y$  be the distance between Ship  $B$  and Lighthouse Rock at time  $t$ , as shown in the figure above.

- Find the distance, in kilometers, between Ship  $A$  and Ship  $B$  when  $x = 4$  km and  $y = 3$  km.
- Find the rate of change, in km/hr, of the distance between the two ships when  $x = 4$  km and  $y = 3$  km.
- Let  $\theta$  be the angle shown in the figure. Find the rate of change of  $\theta$ , in radians per hour, when  $x = 4$  km and  $y = 3$  km.

